

NUB-3172/98-Th

Feb 1998

hep-ph/9802417

Evidence for a Non-Zero Mass in the Dispersion Relation of Transverse Photons in the Stellar Core

Haim Goldberg

Department of Physics

Northeastern University

Boston, MA 02115

Abstract

The electromagnetic two-body interaction energy is modified in the thermal environment of the solar core. The modifications are shown to imply a gross violation of standard solar physics unless the dispersion relation $\omega_t(k)$ for the transverse modes of the photon is altered to include a transverse mass. A non-zero transverse mass $m_t^2 \simeq e^2 n_e / m_e$ is, of course, theoretically predicted to exist; the discussion in this paper implies that such a mass is *required* in order to maintain the success of the standard solar model. In that sense one may claim observational evidence for a non-trivial dispersion relation for transverse photon frequencies in the solar core.

1 Introduction

Electrodynamics (or gluodynamics) in an environment with non-zero temperature or chemical potential is a highly explored topic in modern physics, with references too numerous to list. Historically, one of the first consequences of such considerations [1] was a prediction of long-wavelength oscillations in a neutral plasma with a frequency

$$\omega_{pl} = \sqrt{\frac{e^2 n_e}{m_e}} \quad , \quad (1)$$

where n_e is the electron density. This is an entirely classical effect – essentially a density oscillation of the electron gas which, through coupling via Gauss’s Law, implies a threshold frequency ω_{pl} in the propagation of the longitudinal electric field component $\vec{\nabla} \cdot \vec{E}$. In passing from the classical to the quantum realm, the plasmon mass $\hbar\omega_{pl}$ becomes a zero-momentum energy threshold in a quantized spectrum for collective excitations in the plasma. Experiments verifying the quantum nature of the longitudinal plasmon, in which electrons scattered from thin metallic films show energy losses in multiples of $\hbar\omega_{pl}$, were performed many years ago [2].

As a result of interactions with electrons in the plasma, the dispersion relation $\omega_t(k) \equiv \omega_k$ of the transverse components of the photon also is altered. For small wave vector k , it takes the form [3, 4, 5]

$$\omega_k^2 \simeq k^2 + \omega_{pl}^2 \quad . \quad (2)$$

An extension to the quantum realm of astrophysical significance was suggested by Adams *et al* [4]: with $\hbar\omega_{pl}$ interpreted as a mass for the photon quasiparticle, its decay to a neutrino-antineutrino pair becomes kinematically possible, and could arguably constitute an important mechanism for energy loss in hot dense stellar interiors prior to supernova. It will be some time before this idea can be reliably tested.

The transformation of a quantum-relativistic elementary field particle to a quasiparticle through interaction with the environment is a concept central to many systems of contemporary interest, such as the quark-gluon and the electroweak plasmas, and the theoretical structures required to describe these systems have been extensively developed [6]. It is therefore interesting that there seems as yet to exist little or no

observational support for these ideas. It is the purpose of this work to present in a quantitative manner some evidence, albeit indirect, for the necessity to use a transverse electromagnetic field with the modified dispersion relation (2) in order to describe quantum processes (such as scattering) in the stellar core.

In order to demonstrate that propagators for the quanta of the interacting electromagnetic field require the dispersion relation (2), one necessarily must focus on properties in the environment which involve Feynman graphs (or the equivalent) as contrasted with those solely involving frequencies of classical wave propagation. In this work I examine radiative corrections to the electromagnetic energy density in the core of the sun. I will show that in the thermal environment there are quantum contributions of $\mathcal{O}(e^4)$ to this energy which, for massless transverse photons, are phenomenologically unacceptable. If, on the other hand, the transverse photon indeed behaves like a quasiparticle with mass given by $\hbar\omega_{pl}$, these corrections turn out to be negligible, the energy per particle amounting to no more than $\sim 10^{-5} T_c$, where $T_c \simeq 1.3$ KeV is a typical solar core temperature. It is this argument that I take as (perhaps the first) observational evidence for the existence of transverse quanta with the dispersion relation (2).

2 The Two-Particle Electromagnetic Force in a Thermal Background

Many years ago, Tryon [7] considered the scattering of a charged particle from an external potential in a thermal environment, and calculated $\mathcal{O}(e^2)$ corrections to the cross section due to the presence of background photons. The calculation served as the basis for a speculation in the concluding remarks of that paper that (uncalculated) infrared-finite terms might give rise to interesting phenomena in stellar interiors. In what follows I will calculate a set of thermal corrections to the *static* infrared-finite two-particle Coulomb energy, and assess their effect on stellar physics. These corrections will *not* cancel because of overall charge neutrality.

In the non-relativistic limit, the Hamiltonian describing the interaction of a set of

N charged particles with coordinate vectors $\vec{r}_i$, momentum operators $\vec{p}_i$, charges e_i , and masses m_i with the quantized electromagnetic field $\vec{A}(\vec{r})$ (in Coulomb gauge) is

$$H_{em} = \sum_{i=1}^N \frac{(\vec{p}_i - e\vec{A}(\vec{r}_i))^2}{2m_i} + \frac{1}{2} \sum_{i \neq j=1}^N \frac{e_i e_j}{|\vec{r}_i - \vec{r}_j|} , \quad (3)$$

where the Schrödinger field operator $\vec{A}(\vec{r}_i)$ has the familiar wave number expansion

$$\vec{A}(\vec{r}_i) = \sum_{\vec{k}} \sqrt{\frac{\hbar}{2\omega_k V}} \vec{\epsilon}_{\vec{k}\lambda} \left(a_{\vec{k}\lambda} e^{i\vec{k} \cdot \vec{r}_i} + a_{\vec{k}\lambda}^\dagger e^{-i\vec{k} \cdot \vec{r}_i} \right) , \quad (4)$$

in a real basis for the polarization vector $\vec{\epsilon}_{\vec{k}\lambda}$. In Eq. (3) and in what follows, I will omit factors of c but retain $\hbar$'s so that the quantum properties of various contributions are apparent.

To $\mathcal{O}(e^2)$ and in the non-relativistic limit, the two-body force between pairs of particles $\{i, j\}$ will consist of the (Debye-screened) static Coulomb force and, to $\mathcal{O}(\vec{v}_i \cdot \vec{v}_j / c^2)$, the magnetic Biot-Savart force (which averages to zero). In $\mathcal{O}(e^4)$, aside from simple rescatterings, there are processes, examples of which are shown in the time-ordered graphs of Fig. 1, which make new contributions to the two-body energy.

The thermal environment acts both directly, by allowing photons to be absorbed and re-emitted (as in Figs. 1(b), (c) and (f)), and indirectly, by enhancing the emission and re-absorption of photons in the intermediate state (in all the graphs). In addition, in Coulomb gauge, the diagrams may be classified as containing either one or more *kinetic* vertices from the $\vec{p} \cdot \vec{A}$ terms of Eq. (3), or only *static* contributions from the $\vec{A}^2$ term. Examples of diagrams with kinetic contributions are shown in Figs. 1(d-f). As discussed in the concluding remarks, the results of this paper are fully illustrated by considering only the static contributions: thus, the calculation is limited to an evaluation of Figs. 1(a), (b), and (c). It is also clear from the mass in the denominator of the static term that only the interaction among electrons need be considered.




Figure 1: Various time-ordered graphs contributing to the interaction energy in a thermal environment. The graphs (a), (b), (c) provide the static contributions calculated in this paper.

The contributions from Figs. 1(a), (b), and (c) can be obtained by straightforward calculation. On introducing the expansion (4) into Eq. (3) and making use of second order perturbation theory and a shift to continuum integration, one obtains for Fig. 1(a) (and its partner following the interchange $i \Leftrightarrow j$) the following two-body contribution to the total energy:

$$\begin{aligned} \Delta E_{1a} = & - \left(\frac{e^2}{2m_e} \right)^2 \hbar \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}} \frac{\cos(\vec{k} + \vec{k}') \cdot \vec{r}_{ij}}{\omega_k + \omega_{k'}} \\ & \cdot \sum_{\lambda, \lambda'} \left| \vec{\epsilon}_{\vec{k}\lambda} \cdot \vec{\epsilon}_{\vec{k}'\lambda'} \right|^2 \cdot 2 (n_{\vec{k}\lambda} + 1)(n_{\vec{k}'\lambda'} + 1) , \end{aligned} \quad (5)$$

where

$$n_{\vec{k}\lambda} = \frac{1}{e^{\hbar\omega_k/T} - 1} \quad (6)$$

is the Bose occupation number, and $\vec{r}_{ij} \equiv \vec{r}_i - \vec{r}_j$,

Fig. 1(b) gives a similar result:

$$\begin{aligned} \Delta E_{1b} = & - \left(\frac{e^2}{2m_e} \right)^2 \hbar \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}} \frac{\cos(\vec{k} + \vec{k}') \cdot \vec{r}_{ij}}{\omega_k + \omega_{k'}} \\ & \cdot \sum_{\lambda, \lambda'} \left| \vec{\epsilon}_{\vec{k}\lambda} \cdot \vec{\epsilon}_{\vec{k}'\lambda'} \right|^2 \cdot \left[n_{\vec{k}\lambda} (n_{\vec{k}'\lambda'} + 1) + n_{\vec{k}'\lambda'} (n_{\vec{k}\lambda} + 1) \right] , \end{aligned} \quad (7)$$

while Fig. 1(c) gives

$$\begin{aligned} \Delta E_{1c} = & \left(\frac{e^2}{2m_e} \right)^2 \hbar \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}} \frac{\cos(\vec{k} - \vec{k}') \cdot \vec{r}_{ij}}{\omega_k - \omega_{k'}} \\ & \cdot \sum_{\lambda, \lambda'} \left| \vec{\epsilon}_{\vec{k}\lambda} \cdot \vec{\epsilon}_{\vec{k}'\lambda'} \right|^2 \cdot (n_{\vec{k}\lambda} - n_{\vec{k}'\lambda'}) . \end{aligned} \quad (8)$$

In order to evaluate the thermal contribution in Eqs. (5), (7), and (8), one notes that the integrals are effectively cut off at $k \sim 1/R$, where R is a typical interparticle spacing, so that for massless photons the Bose factors are well approximated by

$$n_{\vec{k}\lambda} \simeq T/\hbar\omega_k \gg 1 . \quad (9)$$

This approximation can be true even if the dispersion relation of ω_k contains an effective mass: in that case, Eq. (9) holds as long as $T \gg \hbar\omega_{pl}$. On substituting (9) into Eqs. (5), (7), and (8), I obtain at $\mathcal{O}(T^2)$

$$\begin{aligned} \Delta E_{1a+1b} = & -4 \left(\frac{e^2}{2m_e} \right)^2 \frac{T^2}{\hbar} \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k^2} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}^2} \frac{\cos(\vec{k} + \vec{k}') \cdot \vec{r}_{ij}}{\omega_k + \omega_{k'}} \\ & \cdot \left(1 + (\hat{k} \cdot \hat{k}')^2 \right) , \end{aligned} \quad (10)$$

while at $\mathcal{O}(T)$

$$\begin{aligned} \Delta E'_{1a+1b} = & -3 \left(\frac{e^2}{2m_e} \right)^2 T \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k^2} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}^2} \cos(\vec{k} + \vec{k}') \cdot \vec{r}_{ij} \\ & \cdot \left(1 + (\hat{k} \cdot \hat{k}')^2 \right) \end{aligned} \quad (11)$$

$$\begin{aligned} \Delta E_{1c} = & - \left(\frac{e^2}{2m_e} \right)^2 T \sum_{i \neq j=1}^N \int \frac{d^3 \vec{k}}{(2\pi)^3 2\omega_k^2} \int \frac{d^3 \vec{k}'}{(2\pi)^3 2\omega_{k'}^2} \cos(\vec{k} - \vec{k}') \cdot \vec{r}_{ij} \\ & \cdot \left(1 + (\hat{k} \cdot \hat{k}')^2 \right) . \end{aligned} \quad (12)$$

In these equations the last factor is a result of performing the polarization sum.

In what follows, I will ignore the $(\hat{k} \cdot \hat{k}')^2$ term in the polarization sum: it can cause no cancellations, and greatly complicates the algebra. With that simplification, the angular integrations can easily be done, with the result

$$\begin{aligned} \Delta E_{1a+1b} = & - \left(\frac{e^2}{4\pi^2} \right)^2 \frac{1}{\hbar} \left(\frac{T}{m_e} \right)^2 \sum_{i \neq j=1}^N \frac{1}{r_{ij}^2} \\ & \cdot \int_0^\infty \frac{k \sin k r_{ij} dk}{\omega_k^2} \int_0^\infty \frac{k' \sin k' r_{ij} dk'}{\omega_{k'}^2} \frac{1}{\omega_k + \omega_{k'}} \end{aligned} \quad (13)$$

$$\begin{aligned} \Delta E'_{1a+1b} + \Delta E_{1c} = & - \left(\frac{e^2}{4\pi^2} \right)^2 \left(\frac{T}{m_e^2} \right) \sum_{i \neq j=1}^N \frac{1}{r_{ij}^2} \\ & \cdot \int_0^\infty \frac{k \sin k r_{ij} dk}{\omega_k^2} \int_0^\infty \frac{k' \sin k' r_{ij} dk'}{\omega_{k'}^2} . \end{aligned} \quad (14)$$

For massless photons, the dispersion relation

$$\omega_k = k \quad (15)$$

holds. In that case, the expressions (13) and (14) (with the appropriate reinstatement of factors of c) become

$$\Delta E_{1a+1b} = - \left(\frac{e^2}{8\pi\hbar c} \right) \left(\frac{T}{m_e c^2} \right)^2 \cdot I \cdot \frac{1}{2} \sum_{i \neq j=1}^N \frac{e^2}{4\pi r_{ij}} \quad (16)$$

$$\Delta E'_{1a+1b} + \Delta E_{1c} = - \left(\frac{e^2}{8\pi} \right) \left(\frac{T}{(m_e c^2)^2} \right) \cdot I' \cdot \frac{1}{2} \sum_{i \neq j=1}^N \frac{e^2}{4\pi r_{ij}^2} , \quad (17)$$

where

$$\begin{aligned} I &= \left(\frac{2}{\pi} \right)^2 \int_0^\infty \frac{\sin x dx}{x} \int_0^\infty \frac{\sin y dy}{y} \frac{1}{x+y} \simeq 0.87 \\ I' &= \left(\frac{2}{\pi} \right)^2 \int_0^\infty \frac{\sin x dx}{x} \int_0^\infty \frac{\sin y dy}{y} = 1 . \end{aligned} \quad (18)$$

It is immediately apparent that (in this massless case) the $\mathcal{O}(T)$ contribution (Eq. (17)) is suppressed over (16) by a factor $\hbar c/TR$, where R is a typical interparticle spacing. For the solar core, with

$$T \simeq 1.3 \text{ KeV}, \quad R \simeq 6 \times 10^9 \text{ cm} \quad ,$$

this is a factor of 10^{-18} . Thus, for the massless case, I focus on Eq. (16), and proceed to estimate it for the solar core environment.

The most important aspect of (16) (and also of (17)) is that it implies an attractive potential between all pairs of particles. This would include the protons as well, but as discussed above, their contribution is suppressed by a large mass factor $(m_e/m_{prot})^2$. With N electrons in a radius R , the leading thermal correction to the total energy is approximately given by

$$\Delta E \simeq -\alpha^2 \left(\frac{T}{m_e} \right)^2 N^2 \frac{\hbar c}{R} \quad , \quad (19)$$

with α the fine structure constant. The relevant measure of this perturbation is its ratio to the thermal energy, NT , which is roughly also the gravitational energy. In natural units ($\hbar = c = 1$), the ratio is seen to be

$$\Delta E/(NT) \simeq -\alpha^2 \left(\frac{T}{m_e} \right) N \frac{1}{m_e R} \quad . \quad (20)$$

For the solar core, $N \simeq 10^{56}$, and $R = 3 \times 10^{17} \text{ KeV}^{-1}$, so that (20) is calculated to be

$$|\Delta E/(NT)| \simeq 10^{29} \quad . \quad (21)$$

This is of course totally unacceptable, and the way out is provided by modifying the dispersion relation for the photon in the thermal bath.

3 Role of the Transverse Mass

For $k \ll m_{pl}$ the effect of the plasma on the photons is a shift in the frequency dispersion relation of from (15) to (2). When the resulting quasiparticle energy spectrum $\hbar\omega_k$ is

introduced in the intermediate states in Eqs. (10) or (13), the result is a Yukawa screening of the long ranges forces. A rough estimate of the modification induced in the energy by (2) can be made as follows:

As noted previously, the integrals (13) and (14) restrict $k \lesssim R^{-1} \ll m_{pl}$. In this case, the transverse photon mass in the stellar core is approximately the plasmon mass (1):

$$\begin{aligned} m_t = \hbar\omega_{pl} &= \sqrt{\frac{4\pi\alpha N_c}{\frac{4\pi}{3}R_c^3 m_e}} \\ &\simeq 400 \text{ eV} , \end{aligned} \quad (22)$$

where R_c is the core radius and N_c is the number of electrons in the core. Since $T/\hbar\omega_{pl} \simeq 3.3$, one may justifiably use the approximation (9) and the simplified form (13) for order-of-magnitude discussion. The integral in (13) may be estimated in the following manner: the introduction of the dispersion relation (2) gives contributions of the form (19), but limited to subvolumes of radius $R_p \leq m_t^{-1}$, with the number of particles N_p in each of these subvolumes given by

$$N_p = N_c (R_p/R_c)^3 . \quad (23)$$

The number of such volumes in the core is N_c/N_p , so that the total contribution to the $\mathcal{O}(T^2)$ energy is (from (19) and (23))

$$\Delta E_{1a+1b} \text{ (screened)} = - \left(\frac{N_c}{N_p} \right) \alpha^2 \left(\frac{T}{m_e} \right)^2 N_p^2 \frac{1}{R_p} \quad (24)$$

$$= \left(\frac{R_p}{R_c} \right)^2 \cdot \Delta E_{1a+1b} \text{ (unscreened)} , \quad (25)$$

The ratio of interaction energy/thermal energy then becomes (omitting $-$ signs)

$$\begin{aligned} \Delta E_{1a+1b} \text{ (screened)} / (N_c T) &= (R_p/R_c)^2 \times 10^{29} \\ &= 7 \times 10^{-35} \times 10^{29} \\ &\simeq 7 \times 10^{-6} , \end{aligned} \quad (26)$$

which renders this effect negligible. A bit more insight is afforded by an algebraic approach: substitute into Eq. (24) the relation (23) and $R_p = m_t^{-1}$ with m_t as given in Eq. (22). The result is

$$\Delta E_{1a+1b} \text{ (screened)}/(N_c T) = \frac{\alpha}{3} \cdot \frac{T}{m_e} . \quad (27)$$

This shows that the suppression depends only on the non-relativistic property of the electrons, which was an assumption of the calculation in the first place.

The $\mathcal{O}(T)$ contribution (14) may be similarly estimated: although it is greatly suppressed over the $\mathcal{O}(T^2)$ term for massless photons, this is not so for quasiparticles. Following the same steps which led to Eq. (27), I find

$$\left(\Delta E'_{1a+1b} + \Delta E_{1c} \right) \text{ (screened)}/(N_c T) \simeq \frac{\alpha}{3} \cdot \frac{m_t}{m_e} , \quad (28)$$

which is again small (of $\mathcal{O}(10^{-6})$), but not greatly suppressed over the $\mathcal{O}(T^2)$ contribution.

4 Summary and Concluding Remarks

(1) In this paper I have calculated corrections to the static two-body Coulomb potential in a thermal photon bath. For massless transverse photons, the interaction energy per particle is many orders of magnitude larger than kT . The screening provided by introducing a photon energy spectrum appropriate to quasiparticles with mass $m_{pl} = \hbar \sqrt{e^2 n_e / m_e}$ is sufficient to suppress this effect in the solar core to negligible significance. In this sense, the argument can be made that quantized transverse quasiparticles are a necessary component for a successful description of physics of the stellar core.

(2) *The Kinetic Terms.* It is not difficult to see that for massless photons and to leading order in temperature T , some of the individual kinetic terms (such as Figs. 1(d) and 1(f)) are infrared divergent. Such divergences will presumably cancel when all contributions are taken into account, in a manner similar to that found by Tryon [7] in his discussion of scattering from an external potential. However, in Coulomb gauge,

even the finite residual part will be distinguished from the static part calculated in this work by having explicit momentum dependence, and hence such contributions can be considered independently of those in this paper. Thus, the results of the calculations in this paper should be sufficient for assessing the necessity of the modified dispersion relation (2).

(3) Because the thermal corrections calculated here, being of $\mathcal{O}(e_i^2 e_j^2)$, are attractive between *all* pairs $\{ij\}$ of particles, the usual Debye screening plays no role in alleviating the energy problem discussed. It is the transverse plasmon mass which is the relevant mitigating factor.

Acknowledgement

This research was supported in part by Grant No. PHY-9722044 from the National Science Foundation.

References

- [1] L. Tonks and I. Langmuir, *Phys. Rev.* **33** (1929) 195.
- [2] C. J. Powell and J. B. Swan, *Phys. Rev.* **115** (1959) 869; **116** (1959) 81.
- [3] V. N. Tsytovich, *Sov. Phys. JETP* **13** (1961) 1249.
- [4] J. B. Adams, M. A. Ruderman, and C.-H. Woo, *Phys. Rev.* **129** (1963) 1383.
- [5] E. Braaten and D. Segel, *Phys. Rev.* **D48** (1993) 1478. This paper presents a calculation of $\omega_t(k)$ over a wide range of wave number k .
- [6] L. Dolan and R. Jackiw, *Phys. Rev.* **D9** (1974) 3320; S. Weinberg, *Phys. Rev.* **D9** (1974) 3357; B. Freedman and L. McLerran, *Phys. Rev.* **D16** (1977) 1130, 1169; J. Kapusta, *Nucl. Phys.* **B148** (1979) 461; D. J. Gross, R. D. Pisarski, and L. G. Yaffe, *Rev. Mod. Phys.* **53** (1981) 43.
- [7] E. P. Tryon, *Phys. Rev. Lett.* **32** (1974) 1139.